

# ON THE SINGULAR SPECTRUM FOR ADIABATIC QUASI-PERIODIC SCHRÖDINGER OPERATORS

M. MARX AND H. NAJAR

**ABSTRACT.** In this paper we study spectral properties of a family of quasi-periodic Schrödinger operators on the real line in the adiabatic limit. We assume that the adiabatic iso-energetic curve has a real branch that is extended along the momentum direction. In the energy intervals where this happens, we obtain an asymptotic formula for the Lyapunov exponent and show that the spectrum is purely singular. This result was conjectured and proved in a particular case by Fedotov and Klopp in [9].

## 1. INTRODUCTION

We consider the following Schrödinger equation

$$(1.1) \quad (H_{z,\varepsilon}\psi)(x) = -\frac{d^2}{dx^2}\psi(x) + [V(x-z) + W(\varepsilon x)]\psi(x) = E\psi(x), \quad x \in \mathbb{R},$$

where  $x \mapsto V(x)$  and  $\zeta \mapsto W(\zeta)$  are periodic, and  $\varepsilon$  is chosen so that the potential  $V(\cdot - z) + W(\varepsilon \cdot)$  be quasi-periodic. Note that in this case, the family of equations (1.1) is ergodic, see [18]; in this case so its spectrum does not depend on  $z$ , see [1]. The operator  $H_{\varphi,\varepsilon}$  can be regarded as an adiabatic perturbation of the periodic operator  $H_0$ :

$$(1.2) \quad H_0 = -\frac{d^2}{dx^2} + V(x).$$

Equation (1.1) is one of the main models of solid state physics. The function  $\psi$  is the wave function of an electron in a crystal with impurities.  $V$  represents the potential of the perfect crystal; as such it is periodic. The potential  $W$  is the perturbation created by impurities. In the semiconductors, this perturbation is slow-varying with respect to the field of the crystal, [21]. It is natural to consider the semi-classical limit.

Let  $\mathcal{E}(\kappa)$  be the dispersion relation associated to  $H_0$ . Consider the *complex* and *real-isoenergy curves*  $\Gamma_{\mathbb{C}}$  and  $\Gamma_{\mathbb{R}}$  defined by

$$(1.3) \quad \Gamma_{\mathbb{C}}(E) = \{(\kappa, \zeta) \in \mathbb{C}^2; \mathcal{E}(\kappa) + W(\zeta) = E\};$$

$$(1.4) \quad \Gamma_{\mathbb{R}}(E) = \{(\kappa, \zeta) \in \mathbb{R}^2; \mathcal{E}(\kappa) + W(\zeta) = E\}.$$

Notice that the iso-energy curves  $\Gamma_{\mathbb{C}}(E)$  and  $\Gamma_{\mathbb{R}}(E)$  are  $2\pi$ -periodic in  $\zeta$  and  $\kappa$  and  $\Gamma_{\mathbb{C}}$  is the Riemann surface uniformizing  $\kappa$ .

The real iso-energy curve has a well known role for adiabatic problems [2]. The adiabatic limit can be regarded as a semi-classical limit and the Hamiltonian  $\mathcal{E}(\kappa) + W(\zeta)$  can be interpreted as a "classical" Hamiltonian corresponding to equation (1.1).

---

1991 *Mathematics Subject Classification.* 34E05, 34E20, 34L05, 34L40.

*Key words and phrases.* quasi-periodic operators, singular spectrum, complex WKB method, Lyapunov exponent.

When  $W$  has in a period exactly one maximum and one minimum, that are non-degenerate, it is proved in [9] that in the energy intervals where the adiabatic iso-energetic curves are extended along the momentum direction, the spectrum is purely singular. This result leads to the following conjecture: in a given interval, if the iso-energy curve has a real branch (a connected component) that is an unbounded vertical curve, then in the adiabatic limit, in this interval, the spectrum is singular. This paper is devoted to prove this conjecture.

Heuristically when the real iso-energy curve is extended along the momentum axis, the quantum states should be extended in momentum and, thus localized in the position space.

**1.1. Results and discussions.** Now, we state our assumptions and results.

**1.1.1. Assumptions on the potentials.** We assume that

**(H1):**  $V$  and  $W$  are periodic,

$$(1.5) \quad V(x+1) = V(x), \quad W(x+2\pi) = W(x).$$

**(H2):**  $V$  is real-valued and locally square-integrable.

**(H3):**  $W$  is real analytic in the strip  $S_Y = \{z \in \mathbb{C} ; |\operatorname{Im} z| < Y\}$ .

We define:

$$(1.6) \quad W_- = \inf_{t \in \mathbb{R}} W(t), \quad W_+ = \sup_{t \in \mathbb{R}} W(t)$$

**1.1.2. Assumptions on the energy region.** To describe the energy regions where we study the spectral properties, we consider the periodic Schrödinger operator  $H_0$  acting on  $L^2(\mathbb{R})$  and defined by (1.2).

**1.1.3. The periodic operator.** The spectrum of (1.2) is absolutely continuous and consists of intervals of the real axis, say  $[E_{2n+1}, E_{2n+2}]$  for  $n \in \mathbb{N}$ , such that  $E_1 < E_2 \leq E_3 < E_4 \dots E_{2n} \leq E_{2n+1} < E_{2n+2} \dots$  and  $E_n \rightarrow +\infty, n \rightarrow +\infty$ . The points  $(E_j)_{j \in \mathbb{N}}$  are the eigenvalues of the self-adjoint operator obtained by considering  $H_0$  defined by (1.2) and acting in  $L^2([0, 1])$  with periodic boundary conditions (see [6, 19]). The intervals  $[E_{2n+1}, E_{2n+2}]$ ,  $n \in \mathbb{N}$ , are the *spectral bands*, and the intervals  $(E_{2n}, E_{2n+1})$ ,  $n \in \mathbb{N}^*$ , the *spectral gaps*. When  $E_{2n} < E_{2n+1}$ , one says that the  $n$ -th gap is *open*; when  $[E_{2n-1}, E_{2n}]$  is separated from the rest of the spectrum by open gaps, the  $n$ -th band is said to be *isolated*. The spectral bands and gaps are represented in figure 1.

**1.1.4. The geometric assumption.** Let us describe the energy region where we study (1.1). We assume that  $J$  is a real compact interval such that:

**(H4a):** For all  $E \in J$ , the connected components of  $\Gamma_{\mathbb{R}}(E)$  have a non empty interior.

**(H4b):** The iso-energy curve  $\Gamma_{\mathbb{R}}(E)$  contains a real branch that is an unbounded vertical curve.

**(H4c):** For all  $E \in J$ , the set  $\mathcal{W}(E) = [E - W_+, E - W_-]$  contains only isolated bands of the periodic operator.




FIGURE 1. Bands, gaps and the quasi-momentum  $k$

$W_+$  and  $W_-$  are defined in (1.6).

## 1.2. Geometric description.

1.2.1. *The set  $W^{-1}(\mathbb{R})$ .* As  $E \in \mathbb{R}$ , the set  $(E - W)^{-1}(\mathbb{R})$  coincide with  $W^{-1}(\mathbb{R})$ . It is  $2\pi$ -periodic. It consists of the real line and of complex branches (curves) which are symmetric with respect to the real line. There are complex branches beginning at the real extrema of  $W$  that do not cross again the real line.

Consider an extremum of  $W$  of order  $n_i$  on the real line, say  $\zeta_i$ . Near  $\zeta_i$ , the set  $W^{-1}(\mathbb{R})$  consists of a real segment, and of  $n_i - 1$  complex curves symmetric with respect to the real axis and intersecting the real axis only on  $\zeta_i$ . The angle between two neighboring curves is equal to  $\frac{\pi}{n_i}$ .

Let  $Y > 0$ . We set  $S_Y = \{-Y \leq \text{Im } \zeta \leq Y\}$ . We assume that  $Y$  is so small that

- $S_Y$  is contained in the domain of analyticity of  $W$ ;
- the set  $W^{-1}(\mathbb{R}) \cap S_Y$  consists of the real line and of the complex lines passing through the real extrema of  $W$ .

An example of subset  $W^{-1}(\mathbb{R})$  is shown in figure 2.

1.2.2. *Notations and description of  $(E - W)^{-1}(\sigma(H_0))$ .* For all  $E \in J$ , we write

$$(E - W)^{-1}(\sigma(H_0)) \cap \mathbb{R} = \bigcup_{k \in \mathbb{Z}} \bigcup_{j=1}^N \{[\varphi_j^-(E), \varphi_j^+(E)] + 2k\pi\},$$

with the following properties:

- $\varphi_1^-(E) < \varphi_1^+(E) < \varphi_2^-(E) < \dots < \varphi_N^-(E) < \varphi_N^+(E), \quad 0 < \varphi_N^+(E) - \varphi_1^-(E) < 2\pi.$




FIGURE 2. A subset of  $W^{-1}(\mathbb{R}) \cap S_Y$

- The connected component  $[\varphi_1^-(E), \varphi_1^+(E)]$  is associated to a connected component of  $\Gamma_{\mathbb{R}}(E)$  which is an unbounded vertical curve.
- We generally define:

$$\varphi_{j+k}^-(E) = \varphi_j^-(E) + 2k\pi, \quad \forall j \in \{1, \dots, N\}, \quad \forall k \in \mathbb{Z}.$$

We set  $\mathcal{B}_j(E) = [\varphi_j^-(E), \varphi_j^+(E)]$ ,  $\mathcal{G}_j(E) = ]\varphi_j^+(E), \varphi_{j+1}^-(E)[$  and  $\mathcal{G}_N(E) = (\varphi_N^+(E), \varphi_1^+(E) + 2\pi) = (\varphi_N^+(E), 2\pi) \cup (2\pi, 2\pi + \varphi_1^-(E))$ . Let  $(\zeta_i^j)_{1 \leq i \leq p_j}$  be the extrema of  $W$  in  $\mathcal{G}_j$ . We recall that  $n_i^j$  is the order of  $\zeta_i^j$ .

We have the following description:

**Lemma 1.1.** *Fix  $[A, B]$  a compact interval of  $\mathbb{R}$ .*

*There exists a finite number  $p$  of real extrema of  $W$  in  $[A, B]$ .*

- *If  $p = 0$ , there exists  $Y > 0$  such that:*

$$(E_0 - W)^{-1}(\mathbb{R}) \cap \{\zeta \in S_Y; \operatorname{Re} \zeta \in [A, B]\} = [A, B].$$

- *For  $p > 0$ . We denote by  $\{\zeta_1, \dots, \zeta_p\}$  the real extremum of  $W$  in  $[A, B]$ . There exist  $Y > 0$  and a sequence  $\{\Sigma_i^1, \dots, \Sigma_i^{n_i-1}\}_{i \in \{1 \dots p\}}$  of disjoint strictly vertical lines of  $\mathbb{C}_+$  starting at  $\zeta_i$  such that:*

$$(E_0 - W)^{-1}(\mathbb{R}) \cap \{\zeta \in S_Y; \operatorname{Re} \zeta \in [A, B]\} = [A, B] \bigcup_{i=1}^p \left( \bigcup_{k=1}^{n_i-1} (\Sigma_i^k \cup \overline{\Sigma_i^k}) \right).$$

1.2.3. *The main result.* The main object of this paper is to prove

**Theorem 1.2.** *We assume (H1)-(H4) are satisfied. For  $\varepsilon > 0$  sufficiently small, for almost all  $z \in \mathbb{R}$ , one has*

$$\sigma(H_{z,\varepsilon}) \cap J \neq \emptyset \quad \text{and} \quad \sigma_{ac}(H_{z,\varepsilon}) \cap J = \emptyset.$$

Here  $\sigma_{ac}(H_{z,\varepsilon})$  is the absolutely continuous spectrum of the family of operators  $(H_{z,\varepsilon})$ .

Using the Ishii-Pastur-Kotani Theorem [4, 18] one can see that the result of Theorem 1.2 is deduced from the positivity of the Lyapunov exponent. We actually recall the following result

**Theorem 1.3.** [4] *Let  $\Theta(E, \varepsilon)$  be the Lyapunov exponent of (1.1). We have*

$$\sigma_{ac}(H_{z,\varepsilon}) = \overline{\{E | \Theta(E) = 0\}}^{ess}.$$

## 2. PERIODIC SCHRÖDINGER OPERATORS

This section is devoted to the study of the periodic Schrödinger operator (1.2) where  $V$  is a 1-periodic, real-valued,  $L^2_{\text{loc}}$ -function. We recall known facts needed on the present paper and we introduce notations. Basic references are [6, 12, 15, 20].

**2.1. Bloch solutions.** Let  $\psi$  be a solution of the equation

$$(2.1) \quad -\frac{d^2}{dx^2}\psi(x, \mathcal{E}) + V(x)\psi(x, \mathcal{E}) = \mathcal{E}\psi(x, \mathcal{E}), \quad x \in \mathbb{R}.$$

satisfying the relation

$$(2.2) \quad \psi(x+1, \mathcal{E}) = \lambda(\mathcal{E})\psi(x, \mathcal{E}).$$

for all  $x \in \mathbb{R}$  and some non-vanishing complex number  $\lambda(\mathcal{E})$  independent of  $x$ . Such a solution exists and is called the *Bloch solution* and  $\lambda(\mathcal{E})$  is called *Floquet multiplier*. We discuss its analytic properties as a function of  $\mathcal{E}$ .

As in section 1.1.2, we denote the spectral bands of the periodic Schrödinger operator by  $[E_{2n+1}, E_{2n+2}]$ ,  $n \in \mathbb{N}$ . Consider  $\mathcal{S}_{\pm}$  two copies of the complex plane  $\mathcal{E} \in \mathbb{C}$  cut along the spectral bands. Paste them together to get a Riemann surface with square root branch points. We denote this Riemann surface by  $\mathcal{S}$ .

One can construct a Bloch solution  $\psi(x, \mathcal{E})$  meromorphic on  $\mathcal{S}$ . It is normalized by the condition  $\psi(1, \mathcal{E}) \equiv 1$ . The poles of this solution are located in the open spectral gaps or at their edges; the closure of each spectral gap contains exactly one pole that, moreover, is simple. It is located either on  $\mathcal{S}_+$  or on  $\mathcal{S}_-$ . The position of the pole is independent of  $x$ .

For  $\mathcal{E} \in \mathcal{S}$ , we denote by  $\widehat{\mathcal{E}}$  the point on  $\mathcal{S}$  different from  $\mathcal{E}$  and having the same projection on  $\mathbb{C}$  as  $\mathcal{E}$ .

We let

$$\widehat{\psi}(x, \mathcal{E}) = \psi(x, \widehat{\mathcal{E}}), \quad \widehat{\mathcal{E}} \in \mathcal{S}.$$

The function  $\widehat{\psi}(x, \mathcal{E})$  is another Bloch solution of (2.1). Except at the edges of the spectrum, the functions  $\psi$  and  $\widehat{\psi}$  are linearly independent solutions of (2.1). In the spectral gaps,  $\psi$  and  $\widehat{\psi}$  are real valued functions of  $x$ , and, on the spectral bands, they differ only by complex conjugation.

**2.2. The Bloch quasi-momentum.** Consider the Bloch solution  $\psi(x, \mathcal{E})$ . The corresponding Floquet multiplier  $\lambda(\mathcal{E})$  is analytic on  $\mathcal{S}$ . Represent it in the form  $\lambda(\mathcal{E}) = \exp(ik(\mathcal{E}))$ . The function  $\mathcal{E} \mapsto k(\mathcal{E})$  is the *Bloch quasi-momentum* of  $H_0$ . Its inverse  $k \mapsto \mathbf{E}(k)$  is the *dispersion relation* of  $H_0$ . A branching point  $\zeta$  is a point where  $k'(\zeta) = 0$ .

Let  $D$  be a simply connected domain containing no branch point of the Bloch quasi-momentum. In  $D$ , one can fix an analytic single-valued branch of  $k$ , say  $k_0$ . All the other single-valued branches of  $k$  that are analytic in  $D$  are related to  $k_0$  by the formulae

$$(2.3) \quad k(\mathcal{E}) = \pm k_0(\mathcal{E}) + 2\pi l, \quad l \in \mathbb{Z}.$$

Consider  $\mathbb{C}_+$ , the upper half plane of the complex plane. On  $\mathbb{C}_+$ , one can fix a single valued analytic branch of the quasi-momentum continuous up to the real line. It can be determined uniquely by the conditions  $\operatorname{Re} k(\mathcal{E} + i0) = 0$  and  $\operatorname{Im} k(\mathcal{E} + i0) > 0$  for  $\mathcal{E} < E_1$ . We call this branch the main branch of the Bloch quasi-momentum and denote it by  $k_p$ .

The function  $k_p$  conformally maps  $\mathbb{C}_+$  onto the first quadrant of the complex plane cut at compact vertical slits starting at the points  $\pi l$ ,  $l \in \mathbb{N}$ . It is monotonically increasing along the spectral zones so that  $[E_{2n-1}, E_{2n}]$ , the  $n$ -th spectral band, is mapped on the interval  $[\pi(n-1), \pi n]$ . Along any open gap,  $\operatorname{Re} k_p(\mathcal{E} + i0)$  is constant, and  $\operatorname{Im} k_p(\mathcal{E} + i0)$  is positive and has only one non-degenerate maximum.

All the branch point of  $k_p$  are of square root type. Let  $E_l$  be a branch point of  $k_p$ . In a sufficiently small neighborhood of  $E_l$ , the function  $k_p$  is analytic in  $\sqrt{E - E_l}$ , and

$$(2.4) \quad k_p(\mathcal{E}) - k_p(E_l) = c_l \sqrt{\mathcal{E} - E_l} + o(\mathcal{E} - E_l) \quad c_l \neq 0.$$

Finally, we note that the main branch can be continued analytically to the complex plane cut along  $(-\infty, E_1]$  and the spectral gaps  $]E_{2n}, E_{2n+1}[$ ,  $n \in \mathbb{N}^*$ , of the periodic operator  $H_0$ .

**2.3. A meromorphic function.** Now let us discuss a function playing an important role in the adiabatic constructions.

In [12], it is shown that, on  $\mathcal{S}$ , there is a meromorphic function  $\omega$  having the following properties:

- the differential  $\Omega = \omega d\mathcal{E}$  is meromorphic; its poles are the points of  $P \cup Q$ , where  $P$  is the set of poles of  $\mathcal{E} \mapsto \psi(x, \mathcal{E})$ , and  $Q$  is the set of zeros of  $k'$ ;
- all the poles of  $\Omega$  are simple;
- if the residue of  $\Omega$  at a point  $p$  is denoted by  $\operatorname{res}_p \Omega$ , one has

$$(2.5) \quad \operatorname{res}_p \Omega = 1, \forall p \in P \setminus Q,$$

$$(2.6) \quad \operatorname{res}_q \Omega = -1/2, \forall q \in Q \setminus P,$$

$$(2.7) \quad \operatorname{res}_r \Omega = 1/2, \forall r \in P \cap Q.$$

- if  $\mathcal{E} \in \mathcal{S}$  projects into a gap, then  $\omega(\mathcal{E}) \in \mathbb{R}$ .

- if  $\mathcal{E} \in \mathcal{S}$  projects inside a band, then  $\overline{\omega(\mathcal{E})} = \omega(\widehat{\mathcal{E}})$ .

2.3.1. *The complex momentum.* It is the main analytic object of the complex WKB method. Let  $\zeta \in S_Y$ . We define  $\kappa$ , in  $\mathcal{D}(W)$  the domain of analyticity of  $W$  by by :

$$(2.8) \quad \kappa(\zeta) = k(E - W(\zeta)).$$

Here,  $k$  is the Bloch quasi-momentum defined in section 2.2. Though  $\kappa$  depends on  $E$ , we omit the  $E$ -dependence. Relation (2.8) translates the properties of  $k$  into properties of  $\kappa$ . Hence,  $\zeta \mapsto \kappa(\zeta)$  is a multi-valued analytic function, and its branch points are related to the branch points of the quasi-momentum by the relations

$$(2.9) \quad E - W(\zeta) = E_l, \quad l = 1, 2, 3, \dots$$

Let  $\zeta_0$  be a branch point of  $\kappa$ . If  $W'(\zeta_0) \neq 0$ , then  $\zeta_0$  is a branch point of square root type.

If  $D \subset \mathcal{D}(W)$  is a simply connected set containing no branch points of  $\kappa$ , we call it *regular*. Let  $\kappa_p$  be a branch of the complex momentum analytic in a regular domain  $D$ . All the other branches that are analytic in  $D$  are described by the following formulae:

$$(2.10) \quad \kappa_m^\pm = \pm \kappa_p + 2\pi m.$$

Here  $\pm$  and  $m \in \mathbb{Z}$  are indexing the branches.

2.3.2. *Index of an interval  $[\varphi_j^-(E), \varphi_j^+(E)]$ .* Fix  $j \in \{1, \dots, N\}$ . Fix a continuous branch  $\kappa_j$  of the complex momentum on  $[\varphi_j^-(E), \varphi_j^+(E)]$ . We define:

$$(2.11) \quad \kappa_j^+ = \kappa_j(\varphi_j^+); \quad \kappa_j^- = \kappa_j(\varphi_j^-); \quad p_j\pi = \kappa_j^+ - \kappa_j^-.$$

$p_j$  is the *index* of  $[\varphi_j^-(E), \varphi_j^+(E)]$  associated to  $\kappa_j$ .

Let us give some properties of  $p_j$ .

**Lemma 2.1.** *Assume that (H4) is satisfied. The indexes  $p_j$  have the following properties:*

- (1) For  $j \in \{1, \dots, N\}$ ,  $p_j \in \{-1, 0, 1\}$ .
- (2)  $\sum_{j=1}^N |p_j| \in 2\mathbb{N}$ .

**Proof** The points  $(E - W)(\varphi_j^-(E))$  and  $(E - W)(\varphi_j^+(E))$  are the ends of a band of  $\sigma(H_0)$ : they are distinct or coincide. If they coincide, i.e. if  $(E - W)(\varphi_j^-(E)) = (E - W)(\varphi_j^+(E))$ , the index  $p_j$  satisfies  $p_j = 0$ . Else, we consider  $k_j$  the branch of the quasi-momentum associated to  $\kappa_j$ . We have that

$$|k_j((E - W)(\varphi_j^-(E))) - k_j((E - W)(\varphi_j^+(E)))| = \pi,$$

see (2.2), and  $|p_j| = 1$ .

Let us prove point (2). We write:

$$\sum_{j=1}^N |p_j| \equiv \sum_{j=1}^N p_j \pmod{2},$$

$$\sum_{j=1}^N p_j = \sum_{j=1}^N \frac{\kappa_j(\varphi_j^+) - \kappa_j(\varphi_j^-)}{\pi} = \frac{\kappa_N(\varphi_N^+) - \kappa_1(\varphi_1^-)}{\pi} + \sum_{j=1}^{N-1} \frac{\kappa_{j+1}(\varphi_{j+1}^-) - \kappa_j(\varphi_j^+)}{\pi}.$$

For  $j \in \{1, \dots, N-1\}$ ,  $(E-W)(\varphi_{j+1}^-(E))$  and  $(E-W)(\varphi_j^+(E))$  are the ends of a same gap and

$$k_{j+1}((E-W)(\varphi_{j+1}^-(E))) \equiv k_j((E-W)(\varphi_j^+(E)))[2\pi].$$

By periodicity,  $(E-W)(\varphi_1^-(E))$  and  $(E-W)(\varphi_N^+(E))$  are the ends of the same gap.

This ends the proof of Lemma 2.1.

If  $p_j \neq 0$ , we say that we cross a band. In this case,  $(E-W)(\varphi_j^-) \neq (E-W)(\varphi_j^+)$  and the associated connected component of the iso-energy curve is unbounded vertically.

We notice that  $p_1 \neq 0$ , and thus  $\sum_{j=1}^N |p_j| > 0$ .

**Remark 2.2.** We can show the determination of  $\kappa$  such that  $p_1 = 1$ .

**2.3.3. Tunneling coefficients.** For  $j \in \{1, \dots, N\}$ , we denote by  $\gamma_j$  a smooth closed curve that goes once around  $[\varphi_j^+(E), \varphi_{j+1}^-(E)]$ . Notice that this curve is the projection of a closed curve on the complex Riemann surface  $\kappa(\zeta) = k(E-W(\zeta))$ . We consider the tunneling actions  $S_j$  given by:

$$(2.12) \quad S_j(E) = i \oint_{\gamma_j} \kappa(\zeta) d\zeta, \quad \forall j \in \{1, \dots, N\},$$

It is straightforward to prove that for  $E \in J$ , each of these actions is real and non-zero and that  $S_j$  is analytic in a complex neighborhood of  $J$  (for analogous statements, we refer to [8, 17]). By definition, we choose the direction of the integration so that all the tunneling actions be positive. We set

$$(2.13) \quad t_j = e^{-\frac{1}{2\varepsilon} S_j},$$

$$T(E, \varepsilon) = e^{-\sum_{j=1}^N S_j(E)/2\varepsilon}.$$

$T$  is exponentially small.

### 3. THE PROOF OF THEOREM 1.2

**3.0.4. Spectral results.** One of the main objects of the spectral theory of quasi-periodic operators is the Lyapunov exponent (for a definition and additional information, see, for example, [4]). The main result of this section is

**Theorem 3.1.** *We assume that the assumptions (H1) – (H4) are satisfied. Then, on the interval  $J$ , for sufficiently small irrational  $\varepsilon/2\pi$ , the Lyapunov exponent  $\Theta(E, \varepsilon)$  of (1.1) is positive and satisfies the asymptotics*

$$(3.1) \quad \Theta(E, \varepsilon) = \frac{\varepsilon}{2\pi} \sum_{j=1}^N \ln \frac{1}{t_j} + o(1) = \frac{1}{4\pi} \sum_{j=1}^N S_j(E) + o(1).$$

This theorem implies that if  $\varepsilon/2\pi$  is sufficiently small and irrational, then, the Lyapunov exponent is positive for all  $E \in J$ .



**3.1. The monodromy matrix and the Lyapunov exponents.** The main object of our study in this subsection is the monodromy matrix for the family of equations (1.1), we define it briefly (we refer the reader to [8, 9]). In this paper, we compute the asymptotics of its Fourier expansion in the adiabatic limit.

3.1.1. *Definition of the monodromy matrix.* Fix  $E \in \mathbb{R}$ . Consider the family of differential equations indexed by  $z \in \mathbb{R}$ ,

$$(3.2) \quad \left( -\frac{d^2}{dx^2} + V(x-z) + W(\varepsilon x) \right) \psi(x) = E\psi(x).$$

**Definition 3.2.** We say that  $(\psi_i)_{i \in \{1,2\}}$  is a consistent basis of solutions to (3.2) if the two functions  $((x, z) \mapsto \psi_i(x, z, E))_{i \in \{1,2\}}$  are a basis of solutions to (3.2) whose Wronskian is independent of  $z$  and that are 1-periodic in  $z$  i.e that satisfy

$$(3.3) \quad \forall x \in \mathbb{R}, \quad \forall z \in \mathbb{R}, \quad \forall i \in \{1,2\}, \quad \psi_i(x, z+1, E) = \psi_i(x, z, E).$$

We refer the reader to [9, 12] about the existence and details on consistent basis of solutions to (3.2).

The functions  $((x, z) \mapsto \psi_i(x+2\pi/\varepsilon, z+2\pi/\varepsilon, E))_{i \in \{1,2\}}$  being also solutions of equation (3.2), we get the relation

$$(3.4) \quad \Psi(x+2\pi/\varepsilon, z+2\pi/\varepsilon, E) = M(z, E)\Psi(x, z, E),$$

where

- $\Psi(x, z, E) = \begin{pmatrix} \psi_1(x, z, E) \\ \psi_2(x, z, E) \end{pmatrix}$ ,
- $M(z, E)$  is a  $2 \times 2$ -matrix with coefficients independent of  $x$ .

The matrix  $M$  is called the *monodromy matrix* associated to the consistent basis  $(\psi_{1,2})$ . We recall the following properties of this matrix.

$$(3.5) \quad \det M(z, E) \equiv 1, \quad M(z+1, E) = M(z, E), \quad \forall z \in \mathbb{R}.$$

The Matrix  $M$  belongs to  $SL(2, \mathbb{R})$  which is known to be isomorph to  $SU(1, 1)$ .

**3.2. The Lyapunov exponents and the monodromy equation.** Consider now a 1-periodic,  $SL(2, \mathbb{C})$ -valued function, say,  $z \mapsto \widetilde{M}$ , and  $h > 0$  irrational. Consider the finite difference equation

$$(3.6) \quad F_{n+1} = \widetilde{M}(z+nh)F_n \quad \forall n \in \mathbb{Z}, \quad F_n \in \mathbb{C}^2.$$

Going from equation (1.1) to the (3.6) is close to the monodromization transformation introduced in [10] to construct Bloch solutions of difference equation. Indeed, it appears that the behavior of solutions of (1.1) for  $x \rightarrow \mp\infty$  repeats the behavior of solutions of the monodromy equation for  $n \rightarrow \mp\infty$ . And it is a well known fact that the spectral properties of the one dimensional Schrödinger

equations can be described in terms of the behavior of its solutions as  $x \rightarrow \mp\infty$ .

The Lyapunov exponent of the finite difference equation (3.6) is

$$(3.7) \quad \theta(\widetilde{M}, h) = \lim_{N \rightarrow +\infty} \frac{1}{N} \log \|P_N(z, h)\|.$$

where the matrix cocycle  $(P_N(z, h))_{N \in \mathbb{N}}$  is defined as

$$(3.8) \quad P_N(z, h) = \widetilde{M}(z + Nh) \cdot \widetilde{M}(z + (N-1)h) \cdots \widetilde{M}(z + h) \cdot \widetilde{M}(z).$$

It is well known that, if  $h$  is irrational, and  $\widetilde{M}$  is sufficiently regular in  $z$ , then the limit (3.7) exists for almost all  $z$  and is independent of  $z$ .

Set  $h \equiv 2\pi/\varepsilon$  [1]. Let  $M$  be the monodromy matrix associated to a consistent basis  $(\psi_{1,2})$ . Consider the *monodromy equation*

$$(3.9) \quad F_{n+1} = M(z + nh, E)F_n \quad \forall n \in \mathbb{Z}, \quad F_n \in \mathbb{C}^2.$$

The Lyapunov exponent of the monodromy equation (3.9) is defined by

$$\theta(E, \varepsilon) = \theta(M(z, E), h).$$

There are several deep relations between equation (3.2) and the monodromy equation (3.9) (see [9, 13]).

We describe only one of them. Recall that  $\Theta(E, \varepsilon)$  is the Lyapunov exponent of the equation (1.1)

We have the following result.

**Theorem 3.3.** ([9]) *Assume  $\varepsilon/2\pi$  is irrational. The Lyapunov exponent  $\Theta(E, \varepsilon)$  and  $\theta(E, \varepsilon)$  are related by the following relation.*

$$(3.10) \quad \Theta(E, \varepsilon) = \frac{\varepsilon}{2\pi} \theta(E, \varepsilon).$$

**3.3. The asymptotics of the monodromy matrix.** As  $W$  and  $V$  are real on the real line, we construct a monodromy matrix of the form

$$(3.11) \quad M(z, E) = \begin{pmatrix} \frac{A(z, E)}{B(\bar{z}, \bar{E})} & \frac{B(z, E)}{A(\bar{z}, \bar{E})} \end{pmatrix}$$

To get (3.11), it suffices to consider a basis of solutions of the form  $(u; \bar{u})$ . For the details on the existence and the construction of such a basis are developed in [8].

The following result gives the asymptotics of  $A$  and  $B$  in the adiabatic case.

**Theorem 3.4.** *Let  $E_0$  be in  $J$ . There exists  $Y > 0$  and  $V_0$ , a neighborhood of  $E_0$ , such that, for sufficiently small  $\varepsilon$ , the family of equations (3.2) has a consistent basis of solutions for which the corresponding monodromy matrix  $M$  is analytic in  $(z, E) \in \{z \in \mathbb{C} ; |Imz| < Y/\varepsilon\} \times V_0$  and has the form (3.11). When  $\varepsilon$  tends to 0, the coefficients  $A$  and  $B$  admit the asymptotics*

$$(3.12) \quad A = A_+(E, \varepsilon)e^{-2iQ_+\pi z}[1 + o(1)], \quad B = B_+(E, \varepsilon)e^{-2iP_+\pi z}[1 + o(1)], \quad 0 < Imz < Y/\varepsilon,$$

$$(3.13) \quad A = A_-(E, \varepsilon)e^{2iQ_-\pi z}[1 + o(1)], \quad B = B_-(E, \varepsilon)e^{2iP_-\pi z}[1 + o(1)], \quad -Y/\varepsilon < Imz < 0.$$

The integers  $P_-, P_+$  and  $Q_-, Q_+ \in \mathbb{Z}$  are specified in section 3. There exists a constant  $C > 1$  such that for  $\varepsilon > 0$  sufficiently small and  $E \in V_0 \cap \mathbb{R}$ , one has

$$(3.14) \quad \frac{1}{C} < T(E, \varepsilon) |A_{\pm}(E, \varepsilon)| < C, \quad \frac{1}{C} < T(E, \varepsilon) |B_{\pm}(E, \varepsilon)| < C,$$

where  $T(E, \varepsilon)$  is defined in (2.13).

For  $Y_1$  and  $Y_2$  such that  $0 < Y_1 < Y_2 < Y$ , there exists  $V = V(Y_1, Y_2)$  a neighborhood of  $E_0$  such that the asymptotics (3.12) and (3.13) for  $A$  and  $B$  are uniform in  $(z, E) \in \{z \in \mathbb{C} ; Y_1/\varepsilon < |\operatorname{Im} z| < Y_2/\varepsilon\} \times V$ .

**Remark 3.5.** The coefficients  $A_+, A_-, B_+$  and  $B_-$  are the leading terms of the asymptotics of the  $Q_{\mp}$ -th and  $P_{\mp}$ -th Fourier coefficients of the monodromy matrix coefficients. From Theorem 3.4, one deduces that, in the strip  $\{-Y < \operatorname{Im} \zeta < Y\}$ , only a few Fourier series terms of the monodromy matrix dominate.

The proof of Theorem 1.2 is given in section 3.

### 3.4. The asymptotics for the Lyapunov exponent.

3.4.1. *The upper bound.* Fix  $P = \max |P_{\mp}|, |Q_{\mp}|$ . The asymptotics (3.12) and (3.13) and estimates (3.14) imply the following estimates for the coefficients of  $M(z, E)$ , the monodromy matrix:

$$(3.15) \quad |A|, |B| \leq C(y_0) \cdot T(E)^{-1} e^{2\pi P y_0/\varepsilon}, \quad \operatorname{Im} z = y_0/\varepsilon,$$

$$(3.16) \quad |A|, |B| \leq C(y_0) \cdot T(E)^{-1} e^{2\pi P y_0/\varepsilon}, \quad \operatorname{Im} z = -y_0/\varepsilon.$$

Here,  $C(y_0)$  is a positive constant independent of  $\varepsilon$ ,  $\operatorname{Re} z$ , and  $E$ . The estimates are valid for sufficiently small  $\varepsilon$ . We recall that  $M$  is analytic and 1-periodic in  $z$ . Equations (3.15), (3.16) and the maximum principle imply that:

$$|A|, |B| \leq 2C(y_0)T(E)^{-1} \exp(2\pi P y_0/\varepsilon), \quad z \in \mathbb{R}.$$

This leads to the following upper bound for the Lyapunov exponent for the matrix cocycle generated by  $M(z, E)$ .

$$(3.17) \quad \theta(E, \varepsilon) \leq \log(T(E)^{-1}) + C + 2\pi P y_0/\varepsilon;$$

where  $C$  is a constant independent of  $E$  and  $\varepsilon$ . Using (3.10) one gets

$$(3.18) \quad \Theta(E, \varepsilon) \leq \frac{\varepsilon}{2\pi} \log(T(E)^{-1}) + \varepsilon C + 2\pi P y_0.$$

3.4.2. *The lower bound.* For  $(M(z, \varepsilon))_{0 < \varepsilon < 1}$  a family of  $SL(2, \mathbb{C})$ -valued, 1-periodic functions of  $z \in \mathbb{C}$  and  $h$  an irrational number, we recall the following result obtained in [10].

**Proposition 3.6.** Fix  $\varepsilon_0 > 0$ . Assume that there exist  $y_0$  and  $y_1$  such that  $0 < y_0 < y_1 < \infty$  and such that, for any  $\varepsilon \in (0, \varepsilon_0)$  one has

- the function  $z \rightarrow M(z, \varepsilon)$  is analytic in the strip  $S = \{z \in \mathbb{C} ; 0 \leq \operatorname{Im} z \leq \frac{y_1}{\varepsilon}\}$ ;

- in the strip  $S_1 = \{z \in \mathbb{C}; \frac{y_0}{\varepsilon} \leq \text{Im}z \leq \frac{y_1}{\varepsilon}\} \subset S$ ,  $M(z, \varepsilon)$  admits the representation

$$(3.19) \quad M(z, \varepsilon) = \lambda(\varepsilon) e^{i2\pi n_0 z} \cdot (M_0(\varepsilon) + M_1(z, \varepsilon));$$

for some constant  $\lambda(\varepsilon)$ , some integer  $n_0$  and a matrix  $M_0(\varepsilon)$ , all of them independent of  $z$ ;

- $M(z, E) = \begin{pmatrix} 1 & 0 \\ \beta(\varepsilon) & \alpha(\varepsilon) \end{pmatrix}$ ;
- there exist constants  $\beta > 0$  and  $\alpha \in (0, 1)$  independent of  $\varepsilon$  such that  $|\alpha(\varepsilon)| \leq \alpha$  and  $|\beta(\varepsilon)| \leq \beta$ ;
- $m(\varepsilon) = \sup_{z \in S_1} \|M_1(z, \varepsilon)\| \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .

Then, there exists  $C > 0$  and  $\varepsilon_1 > 0$  (both depending only on  $y_0, y_1, \alpha, \beta$  and  $\varepsilon \mapsto m(\varepsilon)$ ) such that, if  $0 < \varepsilon < \varepsilon_1$ , one has

$$(3.20) \quad \theta(M(\cdot, \varepsilon), h) > \log |\lambda(\varepsilon)| - Cm(\varepsilon)$$

Proposition 3.6 is used by applying the arguments of [9, 12] to get the lower bound for the Lyapunov exponent.

First for  $\sigma = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  we prove that the matrix  $\sigma M(z, E) \sigma$  completes the assumption of Proposition 3.6.

Let  $y_0$  and  $y_1$  fixed such that  $0 < y_0 < y_1 < Y$ . The asymptotics of the monodromy matrix are uniform for  $z$  in  $S = \{z \in \mathbb{C}; y_0/\varepsilon \leq \text{Im}z \leq y_1/\varepsilon\}$  and  $E \in V_0$ .

Let us assume that  $n = N_N$  in (5.12) is even then the following relations holds

$$(3.21) \quad Q_- = Q_+ + 1, \quad Q_- = P_- + 1 \text{ and } Q_- = P_+$$

For  $E \in V_0 \cap \mathbb{R}$  and  $z \in S$ , Theorem 3.4 implies that

$$\begin{aligned} \overline{A(\bar{z}, E)} &= \overline{A_-(E, \varepsilon)} e^{-2\pi i Q_- z} (1 + o(1)), \quad \frac{A(z, E)}{\overline{A(\bar{z}, E)}} = o(1), \\ \frac{\overline{B(\bar{z}, E)}}{\overline{A(\bar{z}, E)}} &= o(1), \quad \frac{B(z, E)}{\overline{A(\bar{z}, E)}} = c(E)(1 + o(1)), \end{aligned}$$

Where  $c(E)$  is independent of  $z$  and bounded by a constant uniformly in  $\varepsilon$  and  $E$ . So, we have

$$\sigma \cdot M(z, E) \cdot \sigma = \overline{A_-(E, \varepsilon)} e^{-2\pi i Q_- z} \cdot \left( \begin{pmatrix} 1 & 0 \\ c(E) & 0 \end{pmatrix} + o(1) \right).$$

This gives that the matrix-valued function

$$z \mapsto \sigma \cdot M(z, E) \cdot \sigma;$$

satisfies the assumptions of Proposition 3.6.

**Remark 3.7.** When  $n = N_N$  in (5.12) is odd then we get

$$(3.22) \quad Q_+ = Q_- + 1, \quad P_- = Q_+, \quad \text{and } P_+ + 1 = Q_+$$

For  $E \in V_0 \cap \mathbb{R}$  and  $z \in S$ , Theorem 3.4 implies that

$$A(z, E) = A_+(E, \varepsilon) e^{-2\pi i Q_+ z} (1 + o(1)), \quad \frac{\overline{A(\bar{z}, E)}}{A(z, E)} = o(1),$$

$$\frac{\overline{B(\bar{z}, E)}}{A(\bar{z}, E)} = c(E)(1 + o(1)), \quad \frac{B(z, E)}{A(z, E)} = o(1),$$

Where  $c(E)$  is independent of  $z$  and bounded by a constant uniformly in  $\varepsilon$  and  $E$ . So, we have

$$\sigma \cdot M(z, E) \cdot \sigma = A_+(E, \varepsilon) e^{-2\pi i Q_+ z} \cdot \left( \begin{pmatrix} 0 & c(E) \\ 0 & 1 \end{pmatrix} + o(1) \right).$$

The most important properties of the matrix  $M(z, E)$  in Proposition 3.6, is that the bigger eigenvalue is 1 [10].

Using (3.20), we get that the Lyapunov exponent  $\theta(E, \varepsilon)$  of the matrix cocycle associated to  $(M(\cdot, E), h)$ , satisfies the estimates

$$(3.23) \quad \theta(E, \varepsilon) \geq \log |A_-| + o(1).$$

Taking into account (1.2) we get that

$$\Theta(E, \varepsilon) \geq \frac{\varepsilon}{2\pi} \log |A_-| + o(\varepsilon).$$

By (3.14) we get

$$(3.24) \quad \Theta(E, \varepsilon) \geq \frac{\varepsilon}{2\pi} \log(T(E)^{-1}) + O(\varepsilon)$$

3.4.3. *Conclusion.* Now we obtain (3.1), by comparing (3.18) and (3.24). Indeed we see that

$$(3.25) \quad \Theta(E, \varepsilon) = \frac{\varepsilon}{2\pi} \log(T(E)^{-1}) + o(1).$$

The expression of  $T(E)$  given by (3.25) and (3.24) give (3.1) for any  $E \in V_0 \cap \mathbb{R}$ .

Recall that  $V_0 \cap \mathbb{R}$  is an open interval containing  $E_0 \in J$ . The above construction can be carried out for any  $E_0 \in J$ . The end of the proof of Theorem 1.2 follows from the compactness of the interval  $J$ .

#### 4. THE COMPLEX WKB METHOD FOR ADIABATIC PROBLEMS

In this section, following [12, 11, 14], we describe the complex WKB method for adiabatically perturbed periodic Schrödinger equations

$$(4.1) \quad -\frac{d^2}{dx^2} \psi(x) + [V(x) + W(\varepsilon x + \zeta)] \psi(x) = E \psi(x), \quad x \in \mathbb{R}.$$

Here,  $V$  is 1-periodic and real valued,  $\varepsilon$  is a small positive parameter, and the energy  $E$  is complex; one assumes that  $V$  is  $L^2_{\text{loc}}$  and that  $W$  is analytic in a strip in the neighborhood  $S_Y$  of the real line. The parameter  $\zeta$  is an auxiliary complex parameter used to decouple the slow variable  $\zeta = \varepsilon x$  and the fast variable  $x$ . The idea of this method is to study solutions of (4.1) in some domains of the complex plane of  $\zeta$  and, then to recover information on their behavior in  $x \in \mathbb{R}$ . Therefore, for  $D$  a complex domain, one studies solutions satisfying the condition:

$$(4.2) \quad \psi(x+1, \zeta) = \psi(x, \zeta + \varepsilon) \quad \forall \zeta \in D.$$

The aim of the WKB method is to construct solutions to (4.1) satisfying (4.2) and that have simple asymptotic behavior when  $\varepsilon$  tends to 0. This is possible in certain special domains of the complex plane of  $\zeta$ . These domains will depend continuously on  $V, W$  and  $E$ . We shall use these solutions to

compute the monodromy matrix; we consider  $V$  and  $W$  as fixed and construct the WKB objects and solutions in an uniform way for energies in a neighborhood of  $E$ .

**4.1. Standard behavior of consistent solutions.** We start by defining another analytic object central to the complex WKB method, the *canonical Bloch solutions*. Then, we describe the *standard behavior* of the solutions.

**4.1.1. Canonical Bloch solutions.** To describe the asymptotic formulae of the complex WKB method, one needs to construct Bloch solutions to the equation

$$(4.3) \quad -\frac{d^2}{dx^2}\psi(x) + V(x)\psi(x) = \mathcal{E}(\zeta)\psi(x), \quad \mathcal{E}(\zeta) = E - W(\zeta), \quad x \in \mathbb{R};$$

that are moreover analytic in  $\zeta$  on a given regular domain.

Let  $\zeta_0$  be a regular point (i.e.  $\zeta_0$  is not a branch point of  $\kappa$ ). Let  $\mathcal{E}_0 = \mathcal{E}(\zeta_0)$ . Assume that  $\mathcal{E}_0 \notin P \cup Q$ . Let  $U_0$  be a sufficiently small neighborhood of  $\mathcal{E}_0$ , and let  $V_0$  be a neighborhood of  $\zeta_0$  such that  $\mathcal{E}(V_0) \subset U_0$ . In  $U_0$ , we fix a branch of the function  $\sqrt{k'(\mathcal{E})}$  and consider  $\psi_{\pm}(x, \mathcal{E})$ , the two branches of the Bloch solution  $\psi(x, \mathcal{E})$  and  $\Omega_{\pm}$ , the corresponding branches of  $\Omega$  (see section 2.3. For  $\zeta \in V_0$ , we set

$$(4.4) \quad \Psi_{\pm}(x, \zeta) = q(\mathcal{E})e^{\int_{\mathcal{E}_0}^{\mathcal{E}} \Omega_{\pm}} \psi_{\pm}(x, \mathcal{E}), \quad q(\mathcal{E}) = \sqrt{k'(\mathcal{E})}, \quad \mathcal{E} = \mathcal{E}(\zeta).$$

The functions  $\Psi_{\pm}$  are called the *canonical Bloch solutions normalized at the point  $\zeta_0$* .

The properties of the differential  $\Omega$  imply that the solutions  $\Psi_{\pm}$  can be analytically continued from  $V_0$  to any regular domain containing  $V_0$ .

The Wronskian of  $\Psi_{\pm}$  satisfies (see [8])

$$(4.5) \quad w(\Psi_+(\cdot, \zeta), \Psi_-(\cdot, \zeta)) = w(\Psi_+(\cdot, \zeta_0), \Psi_-(\cdot, \zeta_0)) = k'(\mathcal{E}_0)w(\psi_+(\cdot, \zeta_0), \psi_-(\cdot, \zeta_0)).$$

For  $\mathcal{E}_0 \notin Q \cup \{E_l\}$ , the Wronskian  $w(\Psi_+(\cdot, \zeta), \Psi_-(\cdot, \zeta))$  is non-zero.

**4.2. Solutions having standard asymptotic behavior.** Fix  $E = E_0$ . Let  $D$  be a regular domain (i.e.  $D \subset D(W)$ , and simply connected set containing no branch points of  $\kappa$ ). Fix  $\zeta_0 \in D$  so that  $\mathcal{E}(\zeta_0) \notin P \cup Q$ . Let  $\kappa$  be a continuous branch of the complex momentum in  $D$ , and let  $\Psi_{\pm}$  be the canonical Bloch solutions normalized at  $\zeta_0$  defined on  $D$  and indexed so that  $\kappa$  be the quasi-momentum for  $\Psi_+$ .

**Definition 4.1.** Let  $\sigma \in \{+, -\}$ . We say that, in  $D$ , a consistent solution  $f$  has *standard behavior* (or *standard asymptotics*) if

- there exists  $V_0$ , a complex neighborhood of  $E_0$ , and  $X > 0$  such that  $f$  is defined and satisfies (4.1) and (4.2) for any  $(x, \zeta, E) \in (-X, X) \times D \times V_0$ ;
- $f$  is analytic in  $\zeta \in D$  and in  $E \in V_0$ ;

- for any compact set  $K \subset D$ , there exists  $V \subset V_0$ , a neighborhood of  $E_0$ , such that, for  $(x, \zeta, E) \in (-X, X) \times K \times V$ ,  $f$  has the uniform asymptotic

$$(4.6) \quad f = e^{\sigma \frac{i}{\varepsilon} \int_{\zeta_0}^{\zeta} \kappa(u) du} (\Psi_\sigma + o(1)) \text{ as } \varepsilon \text{ tends to } 0;$$

- this asymptotic can be differentiated once in  $x$  without losing its uniformity properties.

We set

$$(4.7) \quad f^*(x, \zeta, E, \varepsilon) = \overline{f(x, \bar{\zeta}, \bar{E}, \varepsilon)}$$

We call  $\zeta_0$  the normalization point for  $f$ . To say that a consistent solution  $f$  has standard behavior, we will use the following notation

$$f \sim \exp \left( \sigma \frac{i}{\varepsilon} \int_{\zeta_0}^{\zeta} \kappa(u) du \right) \Psi_\sigma.$$

### 4.3. Some results on the continuation of asymptotics.

4.3.1. *Description of the Stokes lines near  $[\varphi_1^-(E), \varphi_1^+(E)]$ .* This section is devoted to the description of the Stokes lines under assumption (H4).

4.3.2. *Definition.* The definition of the Stokes lines is fairly standard, [13, 8]. The integral  $\zeta \mapsto \int_{\zeta_0}^{\zeta} \kappa(u) du$  has the same branch points as the complex momentum. Let  $\zeta_0$  be one of them. Consider the curves beginning at  $\zeta_0$ , and described by the equation

$$(4.8) \quad \operatorname{Im} \int_{\zeta_0}^{\zeta} (\kappa(\xi) - \kappa(\zeta_0)) d\xi = 0$$

These curves are the *Stokes lines* beginning at  $\zeta_0$ . According to equation (2.3), the Stokes line definition is independent of the choice of the branch of  $\kappa$ .

Assume that  $W'(\zeta_0) \neq 0$ . Equation (2.4) implies that there are exactly three Stokes lines beginning at  $\zeta_0$ . The angle between any two of them at this point is equal to  $\frac{2\pi}{3}$ . Indeed for  $\zeta$  near  $\zeta_0$ , we have

$$\kappa(\zeta) = \kappa(\zeta_0) + c\sqrt{\zeta - \zeta_0}(1 + o(1)).$$

So

$$\int_{\zeta_0}^{\zeta} (\kappa(\zeta) - \kappa(\zeta_0)) d\zeta = c(\zeta - \zeta_0)^{3/2}(1 + o(1)).$$

4.3.3. *Stokes lines for  $E_0 \in J$ .* We describe the Stokes lines beginning at  $\varphi_1^-(E)$  and  $\varphi_1^+(E)$ . Since  $W$  is real on  $\mathbb{R}$ , the set of the Stokes lines is symmetric with respect to the real line.

First,  $\kappa_1$  is real on the interval  $[\varphi_1^-(E), \varphi_1^+(E)]$ ; therefore this set is a Stokes line starting at  $\varphi_1^-(E)$ . The two other Stokes lines beginning at  $\varphi_1^-(E)$  are symmetric with respect to the real axis. We denote by  $\sigma_1^-(E)$  the Stokes line going downward and by  $\overline{\sigma}_1^-(E)$  its symmetric. Similarly, we denote by  $\sigma_1^+(E)$  and  $\overline{\sigma}_1^+(E)$  the two other Stokes lines starting at  $\varphi_1^+(E)$ ,  $\sigma_1^+(E)$  goes upward. These Stokes lines are represented in figure 3




FIGURE 3. Stokes lines for  $E \in J$

**Lemma 4.2.** *The Stokes lines  $\sigma_1^-(E)$  and  $\sigma_1^+(E)$  satisfy the following properties:*

- *The Stokes lines  $\sigma_1^-(E)$  and  $\sigma_1^+(E)$  stay vertical.*
- *$\sigma_1^-(E)$  and  $\sigma_1^+(E)$  do not intersect one another.*

The proof of this Lemma is similar to the studies done in [9, 14, 16, 17]. We do not give the details.

**4.4. Construction of a consistent basis near  $[\varphi_1^-(E), \varphi_1^+(E)]$ .** We recall this result, proved in [14].

**Proposition 4.3.** [14] *Fix  $E_0 \in J$ , and  $\kappa_1$  a continuous determination of the complex momentum on  $[\varphi_1^-(E), \varphi_1^+(E)]$ . There exists a real number  $Y > 0$ , a complex neighborhood  $\mathcal{V}_1$  of  $E_0$  and a consistent basis  $(f_1, (f_1)^*)$  of solutions of equation (4.1) such that  $f_1$  has the standard asymptotic behavior:*

$$(4.9) \quad f_1 \sim e^{\frac{i}{\varepsilon} \int \tilde{\kappa}_1} \Psi_1^+,$$

and

$$f_1^*(x, \zeta, E) = \overline{f(x, \bar{\zeta}, \bar{E})}.$$

to the left of  $\sigma_1^-(E) \cup \overline{\sigma_1^+(\bar{E})}$  (resp. to the right of  $\sigma_1^+(E) \cup \overline{\sigma_1^-(\bar{E})}$ ). The determination  $\tilde{\kappa}_1$  is the continuation of  $\kappa_1$  through  $\{\zeta \in (\sigma_1^-(E) \cup \overline{\sigma_1^+(\bar{E})}) \cap S_Y ; \operatorname{Im} \kappa_1(\zeta) > 0\}$  (resp. through  $\{\zeta \in (\sigma_1^+(E) \cup \overline{\sigma_1^-(\bar{E})}) \cap S_Y ; \operatorname{Im} \kappa_1(\zeta) < 0\}$ ).

We mimic the analysis done in section 5 of [9]. Precisely, we start by a local construction of the solution  $f$  using canonical domain; then, we apply continuation tools i.e the rectangle Lemma, the adjacent domain principle and the Stokes Lemma.

## 5. THE PROOF OF THEOREM 3.4

The Proof of Theorem 3.4 follows the same ideas as the computations given in section 10.2 of [9]. Below we only give the details for the proof of (3.12) and (3.13).



**5.1. Strategy of the computation.** We now begin with the construction of the consistent basis the monodromy matrix of which we compute. Recall that (H1) – (H4) are satisfied.

In the present section, we construct and study a solution  $f$  of (3.2) satisfying 3.3.

To use the complex WKB method, we perform the following change of variable in (3.2)

$$(5.1) \quad x - z \rightarrow x, \quad \varepsilon z \rightarrow \zeta.$$

Then (3.2) takes the form (4.1). In the new variables, the consistency condition (3.3) becomes (4.2).

Note also that in the new variables, for two solutions to (4.1) to form a consistent basis, in addition to being a basis of consistent solutions, their Wronskian has to be independent of  $\zeta$ .

Consider the basis  $(f_1, f_1^*)$  constructed around  $[\varphi_1^-, \varphi_1^+]$  as in Proposition 4.3. Then the monodromy matrix associated to the basis  $(f_1, f_1^*)$  (defined in section 3.1.1) satisfies:

$$(5.2) \quad \begin{pmatrix} f_1(x, \zeta, E, \varepsilon) \\ f_1^*(x, \zeta, E, \varepsilon) \end{pmatrix} = M(\zeta, E, \varepsilon) \begin{pmatrix} f_1(x, \zeta - 2\pi, E, \varepsilon) \\ f_1^*(x, \zeta - 2\pi, E, \varepsilon) \end{pmatrix}.$$

The aim of this section is the computation of  $M(\zeta, E, \varepsilon)$ . The definition of the monodromy matrix implies that

$$(5.3) \quad A(\zeta) = M_{1,1} = \frac{w(f_1(x + 2\pi, \zeta), f_1^*(x, \zeta))}{w(f_1(x, \zeta), f_1^*(x, \zeta))}, \quad B(\zeta) = M_{12} = \frac{w(f_1(x, \zeta), f_1(x, \zeta + 2\pi))}{w(f_1(x, \zeta), f_1^*(x, \zeta))}.$$

This gives that the monodromy matrix is analytic in  $\zeta$  in the strip  $S_Y$  and in  $E$  in a constant neighborhood of  $E_0$ . By the definition of  $f_1$  we get that  $A$  and  $B$  are  $\varepsilon$ -periodic in  $\zeta \in S_Y$ . This is an immediate consequence of the properties of  $f_1$ .

Therefore, we will compute the Fourier series of  $A$  and  $B$ . The strategy of the computation is based on the ideas of [9] and we first recall some notions presented there. We refer the reader to this paper for more details.

Let  $h$  and  $g$  having a standard asymptotic behavior in regular domains  $D_h$  and  $D_g$  and solutions of (4.1):

$$h \sim e^{\frac{i}{\varepsilon} \int_{\zeta_h}^{\zeta} \kappa_h d\zeta} \psi_h(x, \zeta), \quad \text{and} \quad g \sim e^{\frac{i}{\varepsilon} \int_{\zeta_g}^{\zeta} \kappa_g d\zeta} \psi_g(x, \zeta).$$

Here,  $\kappa_h$ , (resp.  $\kappa_g$ ) is an analytic branch of the complex momentum in  $D_h$  (resp.  $D_g$ ),  $\Psi_h$  (resp.  $\Psi_g$ ) is the canonical Bloch solution defined on  $D_h$  (resp.  $D_g$ ), and  $\zeta_h$  (resp.  $\zeta_g$ ) is the normalization point for  $h$  (resp.  $g$ ).

As the solutions  $h$  and  $g$  satisfy the consistency condition (4.2), their Wronskian is  $\varepsilon$ -periodic in  $\zeta$ .

**5.1.1. Arcs.** We assume that  $D_g \cap D_h$  contains a simply connected domain  $\tilde{D}$ . Let  $\gamma$  be a regular curve going from  $\zeta_g$  to  $\zeta_h$  in the following way: staying in  $D_g$ , it goes from  $\zeta_g$  to some point in  $\tilde{D}$ , then, staying in  $D_h$ , it goes to  $\zeta_h$ . We say that  $\gamma$  is an arc associated to the triple  $h, g$  and  $\tilde{D}$ .

As  $\tilde{D}$  is simply connected, all the arcs associated to the triple,  $h, g$  and  $\tilde{D}$ . As  $\tilde{D}$  is simply connected, all the arcs associated to one and the same triple can naturally be considered as equivalent; we denote them by  $\gamma(h, g, \tilde{D})$ .

We continue  $\kappa_h$  and  $\kappa_g$  analytically along  $\gamma(h, g, \tilde{D})$ . From the properties of  $\kappa$ , we deduce that, for  $V$  a small neighborhood of  $\gamma$ , one has

$$(5.4) \quad \kappa_g(\zeta) = \sigma \kappa_h + 2\pi m, \quad \text{for some } m \in \mathbb{Z}, \quad \text{and} \quad \sigma \in \{1, -1\}.$$

$\sigma = \sigma(h, g, \tilde{D})$  is called the signature of  $\gamma(h, g, \tilde{D})$ , and  $m = m(h, g, \tilde{D})$  the index of  $\gamma(h, g, \tilde{D})$ .

5.1.2. *The meeting domain.* Let  $\tilde{D}$  be as above. We call  $\tilde{D}$  a *meeting domain*, if, in  $\tilde{D}$ , the function  $\text{Im } \kappa_h$  and  $\text{Im } \kappa_g$  do not vanish and are of opposite signs.

Note that, for small values of  $\varepsilon$ , whether  $\zeta \mapsto g(x, \zeta)$  and  $\zeta \mapsto h(x, \zeta)$  increase or decrease is determined by the exponential factor  $e^{\int_{\zeta_g}^{\zeta} \kappa_g d\zeta}$  and  $e^{\int_{\zeta_h}^{\zeta} \kappa_h d\zeta}$ . So, roughly, in a meeting domain, along the lines  $\text{Im } \zeta = \text{Const}$ , the solutions  $h$  and  $g$  increase in opposite directions.

5.1.3. *The action and the amplitude of an arc.* . We call the integral

$$S(h, g, \tilde{D}) = \int_{\gamma(h, g, \tilde{D})} \kappa_g d\zeta,$$

the *action* of the arc  $\gamma = \gamma(h, g, \tilde{D})$ . Clearly, the action takes the same value for equivalent arcs.

Assume that  $\mathcal{E}(\zeta) \notin P \cup Q$  along  $\gamma(h, g, \tilde{D})$ . Consider the function  $q_g = \sqrt{k'(\mathcal{E})}$  and the 1-form  $\Omega_g(\mathcal{E}(\zeta))$  in the definition of  $\Psi_g$ . Continue them analytically along  $\gamma$ . We set

$$(5.5) \quad A(h, g, \gamma) = \left( \frac{q_g}{q_h} \right) \Big|_{\zeta=\zeta_h} e^{\int_{\gamma} \omega_g}.$$

$A$  is called the *amplitude* of the arc  $\gamma$ . The properties of  $\Omega$  imply that the amplitudes of two equivalent arcs  $\gamma(h, g, \tilde{D})$  coincide.

5.2. **Results on the Fourier coefficients.** We recall the following result from [9]

**Proposition 5.1.** *Let  $d = d(h, g)$  be a meeting domain for  $h$  and  $g$ , and  $m = m(h, g, d)$  be the corresponding index. Then*

$$(5.6) \quad w(h, g) = w_m e^{\frac{2\pi i m}{\varepsilon}(\zeta - \zeta_h)} (1 + o(1)); \quad \zeta \in S(d),$$

where  $w_m$  is the constant given by

$$(5.7) \quad w_m = A(h, g, d) e^{\frac{i}{\varepsilon} S(h, g, d)} w(\psi_+(\cdot, \zeta_h), \psi_-(\cdot, \zeta_h)),$$

Here  $\psi_+ = \psi_h$  and  $\psi_-$  is complementary to  $\psi_+$ . The asymptotic (5.6) is uniform in  $\zeta$  and  $E$  when  $\zeta$  stays in a fixed compact of  $S(d)$  and  $E$  in a small enough neighborhood of  $E_0$ .

5.2.1. *The index  $m$ .* Let  $\zeta_0$  be a regular point. consider a regular curve  $\gamma$  going from  $\zeta_0$  to  $\zeta_0 + 2\pi$ . Let  $\kappa$  be a branch of the complex momentum that is continuous on  $\gamma$ . We call the couple  $(\gamma, \kappa)$  a period. Let  $(\gamma_1, \kappa_1)$  and  $(\gamma_2, \kappa_2)$  be two periods. Assume that one can continuously deform  $\gamma_1$  into  $\gamma_2$  without intersecting any branching point. By this we define an analytic continuation of  $\kappa_1$  to  $\gamma_2$ . If the analytic continuation coincide with  $\kappa_2$ , we say that the periods are equivalent.

Consider the branch  $\kappa$  along the curve  $\gamma$  of a period  $(\gamma, \kappa)$ . In a neighborhood of  $\zeta_0$ , the starting point  $\gamma$ , one has

$$\kappa(\zeta + 2\pi) = \sigma\kappa(\zeta) + 2\pi m, \quad \sigma \in \{\pm 1\}, \quad m \in \mathbb{Z}.$$

The numbers  $\sigma = \sigma(\gamma, \kappa)$  and  $m = m(\gamma, \kappa)$  are called respectively the signature and the index of the period  $(\gamma, \kappa)$ . They coincide for equivalent periods.

Recall that

$$\mathcal{G} = \cup_{k \in \mathbb{Z}} \{\cup_{j=1}^N \mathcal{G}_j + 2\pi k\};$$

is the pre-image with respect to  $\mathcal{E}$  of the union of the spectral gaps of  $H_0$ . One has

**Lemma 5.2.** [12] *Let  $(\gamma, \kappa)$  be a period such that  $\gamma$  starts at a point  $\zeta_0 \notin \mathcal{G}$ . Assume that  $\gamma$  intersects  $\mathcal{G}$  exactly  $n$  times ( $n \in \mathbb{N}$ ) and that at all intersection points,  $W' \neq 0$ . Let  $r_1, \dots, r_n$  be the values that  $\text{Re } \kappa$  takes consecutively at these intersection points as  $\zeta$  moves along  $\gamma$  from  $\zeta_0$  to  $\zeta_0 + 2\pi$ . Then,*

$$(5.8) \quad \sigma(\gamma, \kappa) = (-1)^n, \quad m(\gamma, \kappa) = \frac{1}{\pi}(r_n - r_{n-1} + r_{n-2} - \dots + (-1)^{n-1}r_1),$$

$$(5.9) \quad = \frac{(-1)^{n-1}}{\pi}(r_1 - r_2 + \dots + (-1)^{n-1}r_n).$$

### 5.3. The Fourier coefficients.

5.3.1. *For B.* By equation (5.3), we have to compute  $w(f(\cdot, \zeta), f(\cdot, \zeta + 2\pi))$ . With this aim in view, we apply the construction done in section 5.1 with:

$$h(x, \zeta) = f(x, \zeta), \quad g(x, \zeta) = (Tf)(x, \zeta), \quad \text{with} \quad (Tf)(x, \zeta) = f(x, \zeta + 2\pi);$$

and

$$(5.10) \quad D_h = \mathcal{D}, D_g = \mathcal{D} - 2\pi, \quad \zeta_h = \zeta_0, \quad \zeta_g = \zeta_0 - 2\pi,$$

$$(5.11) \quad \forall \zeta \in D_h, \quad \kappa_h(\zeta) = \kappa(\zeta), \quad \forall \zeta \in D_g, \quad \kappa_g(\zeta) = \kappa(\zeta + 2\pi).$$

We will start by

5.3.2. *Above the real line.* We take the meeting domain  $D_0$  as the subdomain of the strip  $0 < \text{Im } \zeta < Y$  between the Stokes lines  $\sigma_1^+ - 2\pi$  and  $\sigma_1^+$ . (In this domain we have  $\text{Im } \kappa_g = -\text{Im } \kappa_h < 0$ ). The arc  $\gamma_0$  connects the point  $\zeta_g$  to  $\zeta_h$ . By (5.10), this defines the period  $(\gamma_0, \kappa_g)$ . Using (5.11), one gets that  $m(f(\cdot, \zeta), f(\cdot, \zeta + 2\pi, D_0)) = m(\gamma_0 + 2\pi, \kappa)$  [9].

We use Lemma 5.2 to compute the index. To do this, we have to compute  $\text{Re } \kappa$  at the intersection of  $\gamma_0 + 2\pi$  and  $\mathcal{G}$ .

As  $\zeta \rightarrow \text{Re } \kappa(\zeta)$  is constant on any connected component  $\mathcal{G}_j$  of  $\mathcal{G}$ . Let us start by defining the index  $\lambda_j^+$  of  $\mathcal{G}_j$ , the result of alternation  $(\dots, +, -, +, -, \dots)$  due to the crossing of  $\mathcal{G}_j$ . We notice that




FIGURE 4. Periods equivalent to  $\gamma_0$  and  $\gamma_1$

$$\lambda_j^+ \in \{-1, 0, 1\}.$$

We set

$$(5.12) \quad N_j = 1 + \sum_{i=1}^{j-1} \left( \sum_{l=1}^{n_i} (o_l - 1) \right).$$

Here  $n_j$  is the number of extremum in  $\mathcal{G}_j$ , and  $o_l$  is the order of the  $l^{\text{th}}$  extremum. The following relations hold

$$(5.13) \quad \lambda_1^+ = \frac{1 + (-1)^{\sum_{l=1}^{n_1} (o_l - 1)}}{2}$$

and for  $2 \leq j \leq N$ ,

$$(5.14) \quad \lambda_j^+ = (-1)^{N_j} \left( \frac{-1 + (-1)^{\sum_{l=1}^{n_j} (o_l - 1)}}{2} \right).$$

Without loss of generality we assume that  $\text{Re } \kappa = 0$  on  $(0, \varphi_1^-)$ . By the above notation, we get that

$$(5.15) \quad m(\gamma_0, \kappa) = (-1)^n \left( p_1 \lambda_1^+ + (p_1 + p_2) \lambda_2^+ + \cdots + \left( \sum_{i=1}^j p_i \right) \lambda_j^+ + \cdots + \left( \sum_{i=1}^N p_i \right) \lambda_N^+ \right) = P_+.$$

Here  $p_i$  is the index of  $\mathcal{B}_i$  and  $n = N_N$ .

Now using (5.3) for  $B$  and Proposition 5.1 we get

$$(5.16) \quad B = A(f, T(f), \tilde{D}_0) e^{\frac{i}{\varepsilon} S(f, T(f), \tilde{D}_0)} \cdot e^{\frac{-2i\pi P_+ \zeta_0}{\varepsilon}} = B_+(E, \varepsilon) \cdot e^{-\frac{i}{2\pi\varepsilon} P_+ \zeta_0}, \quad T(f)(x, \zeta) = f(x, \zeta + 2\pi).$$

### 5.3.3. Below the real line.

Below the real line, we take the domain of the strip  $-Y < \text{Im } \zeta < 0$  located between the stokes line  $\sigma_1^- - 2\pi$  and  $\sigma_1^-$  as a regular domain; which we denote by  $\tilde{D}_0$ . We set  $\tilde{\gamma}_0 = \gamma(f, T(f), \tilde{D}_1)$ , then it defines a period, and so  $m(f, T(f), \tilde{D}_1) = m(\tilde{\gamma}_0 + 2\pi, \kappa)$ . The curve defining a period equivalent to  $(\tilde{\gamma}_0 + 2\pi, \kappa)$  is represented in figure 4.

The computation of the index  $\lambda_j^-$  of  $\mathcal{G}_j$  gives in this case that

$$\lambda_1^- = \frac{-1 + (-1)^{\sum_{i=1}^{n_1} (o_i - 1)}}{2},$$

and

$$\lambda_j^- = (-1)^{N_j} \left( \frac{-1 + (-1)^{\sum_{l=1}^{n_j} (o_l - 1)}}{2} \right).$$

By this notation we get that

$$(5.17) \quad m(\gamma_0, \kappa) = (-1)^n \left( p_1 \lambda_1^- + (p_1 + p_2) \lambda_2^- + \cdots + \left( \sum_{i=1}^j p_i \right) \lambda_j^- + \cdots + \left( \sum_{i=1}^N p_i \right) \lambda_N^- \right) = P_-.$$

Using (5.3) for  $B$  and Proposition 5.1 we get

$$(5.18) \quad B_{P_-} = A(f, T(f), \tilde{D}_1) e^{\frac{i}{\varepsilon}(S(f, T(f), \tilde{D}_1) - 2\pi P_- \zeta_0)} = B_-(E, \varepsilon) \cdot e^{\frac{i}{\varepsilon} 2\pi P_- \zeta_0}, \quad (Tf)(x, \zeta) = f(x, \zeta + 2\pi).$$

5.4. **For  $A$ .** For the computation of  $A$  using equation (5.3), we have to compute  $w(f^*(\cdot, \zeta), f(\cdot, \zeta + 2\pi))$ .

It suffices to apply the method presented in section 5.1 with:

$$h(x, \zeta) = f^*(x, \zeta), \quad g(x, \zeta) = (Tf)(x, \zeta), \quad \text{with} \quad (Tf)(x, \zeta) = f(x, \zeta + 2\pi);$$

and

$$(5.19) \quad D_h = \mathcal{D}^*, \quad D_g = \mathcal{D} - 2\pi, \quad \zeta_h = \zeta_0, \quad \zeta_g = \zeta_0 - 2\pi,$$

$$(5.20) \quad \forall \zeta \in D_h, \quad \kappa_h(\zeta) = -\overline{\kappa}(\overline{\zeta}), \quad \forall \zeta \in D_g, \quad \kappa_g(\zeta) = \kappa(\zeta + 2\pi).$$

5.4.1. *Above the real line.* In this case,  $\tilde{D}_0$ , the meeting domain, is the subdomain of the strip  $\{0 < \text{Im} \zeta < Y\}$  located between the lines  $\sigma_1^+ - 2\pi$  and  $\overline{\sigma_1^-}$ , the symmetric to  $\sigma_1^-$  with respect to  $\mathbb{R}$ . The arc  $\gamma(f^*, T(f), \tilde{D}_0)$  defines a period  $(\tilde{\gamma}_0, \kappa_g)$ , in figure 5 we represent the curve  $\tilde{\gamma}_0 + 2\pi$ .

Similarly to the computation of  $B$ , we define the index  $\beta_j^+$  of  $\mathcal{G}_j$  and in this case, we have:

$$\beta_1^+ = \frac{-1 + (-1)^{\sum_{i=1}^{n_1} (o_i - 1)}}{2},$$

and

$$\beta_j^+ = (-1)^{N_j} \left( \frac{-1 + (-1)^{\sum_{i=1}^{l=n_j} (o_i - 1)}}{2} \right).$$

By this notation we get that

$$(5.21) \quad m(\gamma_0, \kappa) = (-1)^n \left( p_1 \beta_1^+ + (p_1 + p_2) \beta_2^+ + \cdots + \left( \sum_{i=1}^j p_i \right) \beta_j^+ + \cdots + \left( \sum_{i=1}^N p_i \right) \beta_N^+ \right) = Q_+.$$

Using (5.3) for  $A$  and Proposition 5.1 we get that

$$m(f^*, Tf, \tilde{D}_0) = m(\tilde{\gamma}_0, \kappa) = Q_+,$$

and

$$(5.22) \quad A = A(f^*, Tf, \tilde{D}_0) e^{\frac{i}{\varepsilon}(S(f^*, Tf, \tilde{D}_0) - 2\pi Q_+ \zeta_0)} = A_+(E, \varepsilon) \cdot e^{-\frac{i}{\varepsilon} 2\pi Q_+ \zeta_0}, \quad (Tf)(x, \zeta) = f(x, \zeta + 2\pi).$$




FIGURE 5. Periods equivalent to  $\tilde{\gamma}_0$  and  $\tilde{\gamma}_1$

5.4.2. *Below the real line.* In this case,  $\tilde{D}_1$ , the meeting domain, is the subdomain of the band  $\{-Y < \text{Im}\zeta < 0\}$  located between the lines  $\overline{\sigma_1^+}$ , symmetric of  $\sigma_1^+$  with respect to  $\mathbb{R}$ , and  $\sigma_1^- - 2\pi$ . The arc  $\gamma(h, g, \tilde{D}_1)$  defines a period  $(\tilde{\gamma}_1, \kappa_g)$ ; the curve  $\tilde{\gamma}_1 + 2\pi$  is represented in figure 5. One obtains that

$$(5.23) \quad \beta_1^- = \frac{1 + (-1)^{\sum_{l=1}^{n_1} (o_l - 1)}}{2},$$

and for  $2 \leq j \leq N$ ,

$$(5.24) \quad \beta_j^- = (-1)^{N_j} \left( \frac{-1 + (-1)^{\sum_{l=1}^{n_j} (o_l - 1)}}{2} \right).$$

So by the above notation we get that

$$(5.25) \quad m(\gamma_0, \kappa) = (-1)^n \left( p_1 \beta_1^- + (p_1 + p_2) \beta_2^- + \cdots + \left( \sum_{i=1}^j p_i \right) \beta_j^- + \cdots + \left( \sum_{i=1}^N p_i \right) \beta_N^- \right) = Q_-$$

One obtains that  $m(f^*, T(f), \tilde{D}_1) = m(\tilde{\gamma}_1 + 2\pi, \kappa) = Q_-$ , and

$$(5.26) \quad A = A(f^*, T(f), \tilde{D}_1) e^{\frac{i}{\varepsilon} (S(f^*, T(f), \tilde{D}_1) - 2\pi Q_- - \zeta_0)} = A_-(E, \varepsilon) \cdot e^{\frac{i}{\varepsilon} 2\pi Q_- - \zeta_0} ; \quad (Tf)(x, \zeta) = f(x, \zeta + 2\pi).$$

**Acknowledgements.** *The authors would like to thank F. Klopp for proposing this problem and many valuable comments and remarks. H.N thanks M. Dimassi for suggesting some interesting references*

## REFERENCES

- [1] J. Avron and B. Simon. Almost periodic Schrödinger operators, II, the Integrated density of states *Duke Mathematical Journal*, 50:369-391, 1983.
- [2] V. S. Buslaev and A. Fedotov: *Bloch solutions for difference equations*. Algebra i Analiz **(7)** p74-122 (1995)
- [3] R. Carmona and J. Lacroix: *Spectral Theory of Random Schrödinger Operators* Birkhäuser Boston. Basel.Berlin.
- [4] H.L Cycon, R.G. Froese, W. Kirsch and B. Simon *Schrödinger operators*. Springer Verlag, Berlin, 1987.
- [5] V. S. Buslaev and A. Fedotov: *The complex WKB method for Harper's equation* St Petersburg Math Journal, **(95)** 6: p495-517 (1995).
- [6] R. Eastham: *The spectral theory of periodic differential operators* Scottish Academic Press, Edinburgh (1973)
- [7] E. Sorets and T. Spencer: *Positive Lyapunov exponents for Schrödinger operators with quasi-periodic potentials*. Comm. Math. Phys. **(142)** (3) p 543-566 (1991)
- [8] A. Fedotov and F. Klopp: *A complex method for adiabatic problems* Asymp. Anal. (2001) **(27)** p 219-264.
- [9] A. Fedotov and F. Klopp: *On the singular Spectrum for Adiabatic Quasi-periodic Schrödinger Operators in the real Line*. Ann. henri Poincaré (2005) **5** p1-50.
- [10] A. Fedotov and F. Klopp: *Anderson transitions for a family of almost periodic Schrödinger equations in the adiabatic case* Comm. Math. Phys. (2002) **(227)** (1) p1-92.
- [11] A. Fedotov and F. Klopp: *Geometric tools of the adiabatic complex WKB method* Asymp. Anal. (2004)
- [12] A. Fedotov and F. Klopp: *On the absolutely continuous spectrum of one dimensional q uasi-periodic schrödinger operators in the adiabatic limit*. Trans. Amer. Math. Soc (2005).
- [13] M. V. Fedoryuk: *Asymptotic Analysis*, Springer Verlag, Berlin; (1993).

- [14] F. Klopp and M. Marx, *Resonances for slowly varying perturbations of one-dimensional periodic Schrödinger operators*, Expos IV. Actes du Sminaire quations aux Drives Partielles de l'Ecole Polytechnique (2005-2006), Exp. No. IV, 18 pp.
- [15] H. MacKean and P. van Moerbeke: *The spectrum of Hill's equation* Inven. Math. **(30)** p 217-274 (1975).
- [16] M. Marx: *Etude de perturbations adiabatiques de l'équation de Schrödinger périodique* Thèse, Université Paris XIII", (2004).
- [17] M. Marx: *On the eigenvalues for slowly varying perturbations of a periodic Schrödinger operator* Asymp. Ana. (2005) **(48)**, N 4, p 295-357 (2006)
- [18] L. Pastur and A. Figotin: *Spectra of Random and Almost-Periodic Operators* Springer-Verlag.
- [19] M. Reed and B.Simon: *Methods of Modern Mathematical vol IV: Analysis of Operators* Academic Press, New
- [20] E.C Titchmarsh: *Eigenfunction expansions associated with second-order differential equations Part II*", Clarendon Press, Oxford (1958). York, 1978.
- [21] J.M Ziman, *Principles of the Theory of Solids*, Cambridge at the University Press, 1972.

(M. Marx) LAGA, U.M.R. 7539 C.N.R.S, INSTITUT GALILÉE, UNIVERSITÉ DE PARIS-NORD, 99 AVENUE J.-B. CLÉMENT, F-93430 VILLETANEUSE, FRANCE

*E-mail address:* [marx@math.univ-paris13.fr](mailto:marx@math.univ-paris13.fr)

(Hatem Najar) DPARTEMENT DE MATHMATIQUES I.S.M.A.I. KAIROUAN, RUE ASSED IBN ELFOURAT 3100 KAIROUAN, TUNISIA

*E-mail address:* [hatem.najar@ipeim.rnu.tn](mailto:hatem.najar@ipeim.rnu.tn)